

Transfer of Care

Inclusion Criteria

- ➤ CTAS 3-5
- Stable vitals
 - HR between 50-100 BPM
 - Respiratory Rate ≤24
 - Systolic BP ≥100 mmHg
 - O2 sat ≥94% on room air
 - Blood Glucose ≥4 mmol/L (if indicated)
 - Afebrile
- GCS 15 or normal cognitive state
- Patient is >14 and <75 years of age or is with a responsible adult or guardian
- Able to ambulate independently or with minimal assistance

Pertinent Negatives

- ➤ **NO** seizure activity within last 24 hours
- > NOT visibly intoxicated
- ➤ Patient is **NOT** experiencing an acute psychiatric emergency
- NO ongoing medical treatment (by Paramedics)
- Does NOT require active monitoring
- ➤ **NO** medications given by Paramedics, other than:
 - Acetaminophen
 - Ibuprofen
 - Ketorolac

| with minimal assistar ➤ Able to sit in chair or ➤ Able to follow direction | nce wheelchair | lotorolad |
|--|-------------------|-----------|
| Most Recent Vital Signs If treatment was provided, vital signs must be at least 15 minutes post treatment. | | |
| Time: | BP: | HR: |
| RR: Temp: | SAT: | BG: |
| | | |
| Pre-Treatment – if applicable: Pre-treatment vital signs must be recorded if medication has been administered by Paramedics: | | |
| Medication Name/Dose: | | |
| Time of Administration: | | |
| Time: | BP: | HR: |
| RR: Temp: | SAT: | BG: |
| Paramedic Signature: | | |

Version 1.4 March of 2023